

# UNCLASSIFIED

AD NUMBER
AD404792
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; FEB 1963. Other requests shall be referred to US Army Biological Laboratories, Fort Detrick, MD 21701.
AUTHORITY
SMUFD, d/a ltr, 4 Feb 1972

THIS PAGE IS UNCLASSIFIED

**UNCLASSIFIED**

**AD 404 792**

---

**DEFENSE DOCUMENTATION CENTER**

**FOR**

**SCIENTIFIC AND TECHNICAL INFORMATION**

**CAMERON STATION, ALEXANDRIA, VIRGINIA**



**UNCLASSIFIED**

**NOTICE:** When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

404 792

63-3-5


CATALOGED BY ASTIA  
AS AD NO. 404792

**A CHANGE OF THE PROPERTIES  
OF THE Q-FEVER CAUSATIVE AGENT**

**TRANSLATION NO.**

**747**

**FEBRUARY 1963**



**U.S. ARMY BIOLOGICAL LABORATORIES  
FORT DETRICK, FREDERICK, MARYLAND**

ASTIA AVAILABILITY NOTICE

Qualified requestors may obtain copies of this document from ASTIA.

This publication has been translated from the open literature and is available to the general public. Non-DOD agencies may purchase this publication from the Office of Technical Services, U. S. Department of Commerce, Washington 25, D. C.

Journal of Microbiology, Epidemiology and Immunobiology (Russian Journal No. 1, 1963, pp. 152). Translated by Sp/6 Charles T. Ostertag, Jr.

A Change of the Properties of the Q-fever Causative Agent, Communication I. Biological Properties of Variants D and B-D-1 of the Grit Strain of Rickettsia burnetii (Author's Summary). Kravchenko, A.T.; Yakovlev, A.I.; Gudima, O.S.; and Polozov, A.I. (Submitted to Editorial Office 16 July 1961).

Results are presented of the experimental study of a mildly virulent variant of the Grit strain of Rickettsia burnetii. The strain was obtained in 1954 from the Gamaleya Institute of Epidemiology and Microbiology of the USSR Academy of Medical Sciences after 44 passages in chick embryos. They supported the strain further by passages in the vitelline membrane of chick embryos. They didn't study the virulence of a culture of the 44th - 52nd passage in chick embryos, and when pigs were infected with large doses of a culture of the 53rd passage in chick embryos, a febrile reaction was absent in the animals, but 21 days after the administration, specific complement-fixing antibodies were detected in the blood. This variant of Grit strain was designated as Grit-D.

In comparative tests it was established that in guinea pigs infected with the original Grit strain, febrile forms of the infection were sometimes encountered during infection with small doses of rickettsias. Infiltrates appeared at the site of administration of the causative agent, as a rule simultaneously with an increase in temperature. During contamination with a sufficiently concentrated culture (dilution  $10^{-1}$  -  $10^{-4}$ ), in several cases there were observed necrotization and ulceration at the site of administration of the rickettsias. Variant D regularly caused a febrile reaction in animals just with the administration of the concentrated material (dilution of the culture  $10^{-2}$  -  $10^{-3}$ ). Along with this, in many of the animals the fever confined itself to an increase of temperature up to  $39.6 - 39.8^{\circ}$  in the course of 2-3 days. In spite of the sharp lowering of the causative agent's ability to cause a febrile illness in guinea pigs, the ability to form infiltrates was preserved (in individual animals a local reaction was observed with the administration of a culture up to a dilution of  $10^{-7}$  -  $10^{-8}$ ). However, their intensity and duration was considerably less than in animals infected with the original Grit strain. The titers of complement-fixing antibodies in guinea pigs infected with variant D and the Grit strain are in direct relation to the dose of rickettsias administered, but with one and the same doses they were considerably lower in animals infected with variant D. The analogous rules were also established in comparative experiments for determining the virulence of variant D and the Grit strain during the intraperitoneal, intramuscular, and intracutaneous methods of infecting guinea pigs.

With the aim of determining the stability of the reduction of pathogenic properties, variant D was passed 5 times through the testicle

of guinea pigs and one time each through two species of Ixodoidea ticks - Lipicephalus sanguineus and Haemaphysalis concinna.

In guinea pigs, infected both with the original variant D and with variants obtained after passage, the duration, incubation period, height and duration of fever, the presence and degree of intensity of infiltrates at the site of administration of the rickettsias, and also their reverse development turned out to be identical.

Finally Rickettsia burnetii (variant D) were subjected to the action of ionizing irradiation. An aqueous solution of radioactive phosphorus ( $\text{Na}_2\text{HP}^{32}\text{O}_4$ ) served as the source of radiation. A study of the variant, obtained after a 4-multiple irradiation with doses of 500, 1200, 1800 and 2520 mc for 1 liter, and 22 subsequent passages in chick embryos (BD-1), showed that its virulence became lower than in the original variant D. However, after two additional passages in chick embryos without irradiation, its virulence was restored.

Observations showed that variant D, based on its biological properties, doesn't differ from variant M which is spontaneously separated from the Grit strain described earlier by Zdrodovskiy and Genig. The similar biological properties of the specified variants and the community of the source where they were obtained (Museum of Rickettsial Strains of the Gamaleya Institute of Epidemiology and Microbiology of the USSR Academy of Medical Sciences) allow us to maintain that variant D is identical to variant M.